


ENVIRONMENTAL SENSITIVITY INDEX MAP

89°30'00"




89°30'00"

SHORELINE HABITATS (ESI)



- 1B EXPOSED, SOLID MAN-MADE STRUCTURES
- 2A EXPOSED WAVE-CUT PLATFORMS IN CLAY OR MUD
- 2B EXPOSED SCARPS AND STEEP SLOPES IN CLAY OR MUD
- 3A FINE- TO MEDIUM-GRAINED SAND BEACHES
- 3B SCARPS AND STEEP SLOPES IN SAND
- 4 COARSE-GRAINED SAND BEACHES
- 5 MIXED SAND AND GRAVEL (SHELL) BEACHES
- 6A GRAVEL BEACHES
- 6B RIPRAP
- 7 EXPOSED TIDAL FLATS
- 8A SHELTERED SCARPS IN CLAY OR MUD
- 8B SHELTERED, SOLID MAN-MADE STRUCTURES
- 8C SHELTERED RIPRAP
- 8E PEAT
- 9A SHELTERED TIDAL FLATS
- 9B VEGETATED LOW BANKS
- 10A SALT-AND BRACKISH-WATER MARSHES
- 10B FRESHWATER MARSHES
- 10C SWAMPS
- 10D SCRUB-SHRUB WETLANDS, INCLUDING BLACK MANGROVES



SCALE 1:50000



Not For Navigation
 Published: December 2013
 Published at Seattle, Washington
 National Oceanic and Atmospheric Administration
 National Ocean Service
 Office of Response and Restoration
 Emergency Response Division



Louisiana: ESIMAP 121 (cont.)

BIOLOGICAL RESOURCES: (cont.)

MARINE MAMMAL:

| RAR# | Species | S F Conc. | J | F | M | A | M | J | J | A | S | O | N | D | Mating | Calving | Pupping | Molting |
|------|---------------------|----------------------|---|---|---|---|---|---|---|---|---|---|---|---|--------|---------|---------|---------|
| 303 | Bottlenose dolphin | VERY ABUNDANT | X | X | X | X | X | X | X | X | X | X | X | X | - | - | - | - |
| 304 | West Indian manatee | E E RARE TO UNCOMMON | X | X | X | X | X | X | X | X | X | X | X | - | - | - | - | |

REPTILE:

| RAR# | Species | S F Conc. | J | F | M | A | M | J | J | A | S | O | N | D | Nesting | Hatching | Interesting | Juveniles | Adults |
|------|--------------------------|----------------|---|---|---|---|---|---|---|---|---|---|---|---|---------|----------|-------------|-----------|--------|
| 177 | Diamondback terrapin | | | | X | X | X | X | X | X | X | | | - | - | - | - | - | |
| 282 | Kemp's ridley sea turtle | E E ABUNDANT | | | X | X | X | X | X | | | | | - | - | - | APR-SEP | - | |
| 284 | Green sea turtle | T T OCCASIONAL | | | X | X | X | X | X | X | X | | | - | - | - | MAR-NOV | MAR-NOV | |
| | Hawksbill sea turtle | E E VERY RARE | | | X | X | X | X | X | X | | | | - | - | - | MAR-OCT | - | |
| | Leatherback sea turtle | E E RARE | X | X | X | X | X | X | X | X | X | X | X | - | - | - | JAN-DEC | JAN-DEC | |
| | Loggerhead sea turtle | T T COMMON | X | X | X | X | X | X | X | X | | | | - | - | - | MAR-NOV | MAR-NOV | |

HUMAN USE RESOURCES:

CRITICAL HABITAT:

| HUN# | Name | Contact | Phone |
|------|--------------------------------|-------------|-------|
| 75 | GULF STURGEON CRITICAL HABITAT | USEFWS/NMFS | |

HELIPORT:

| HUN# | Name | Contact | Phone |
|------|-----------|---------|-------|
| 727 | JACK KENT | | |

MANAGEMENT AREA:

| HUN# | Name | Contact | Phone |
|------|--------------------|------------------------------------|-------|
| 771 | BILOXI WMA | LDWF | |
| 775 | OYSTER LEASE | LDWF - OYSTER LEASE SURVEY SECTION | |
| 779 | PUBLIC OYSTER AREA | LDWF | |

Biological information shown on the maps represents known concentration areas or occurrences, but does not necessarily represent the full distribution or range of each species. The LDWF-LNHP provided information for some of the federally and state listed species and species of conservation concern for display in the ESI atlas and accompanying digital data in 2013. The available LNHP data sets are to be used for oil spill response and spill response planning only. These data represent existing information known to the LNHP at the time of the request and should never be substituted for consultation with the LNHP. The more spatially generalized 2011 polygonal waterbird colony data was provided by LNHP and the more spatially specific 2006 point waterbird colony data was provided by BTNEP. The display of these two data sets does not imply that EITHER or BOTH sets of polygons and/or points (especially if counts are aggregated) reflect current nest locations OR counts, but rather are to be used as a guide for what species could be present.